

QUITE COMPLETE REAL CLOSED FIELDS

SAHARON SHELAH

ABSTRACT. We prove that any ordered field can be extended to one for which every decreasing sequence of bounded closed intervals, of any length, has a nonempty intersection; equivalently, there are no Dedekind cuts with equal cofinality from both sides. Here we strengthen the results from the published version.

1. INTRODUCTION

Laszlo Csirmaz raised the question of the existence of nonarchimedean ordered fields with the following completeness property: any decreasing sequence of closed bounded intervals, of any ordinal length, has nonempty intersection. We will refer to such fields as *symmetrically complete* for reasons indicated below.

Theorem 1.1. *Let K be an arbitrary ordered field. Then there is a symmetrically complete real closed field containing K .*

The construction shows that there is even a “symmetric-closure” in a natural sense, and that the cardinality may be taken to be at most $2^{|K|^{++\aleph_1}}$.

I thank the referee for rewriting the paper.

In September 2005 (after the paper appeared), lecturing on it in the Rutgers logic seminar without details, Cherlin asked where $\kappa \geq d(K)$ was used in the argument. Checking the proof, it appears that this was used (in the published version) but we can improve the result: the completion takes $\leq \omega_2 + 1$ steps instead of a bound depending on K .

Note that by [2], consistently with ZFC is (i.e., after forcing extension) there is an $\aleph_1$ -saturated ultrapower of the field of the real (hence a real closed field) which is Scott complete. Also compared to the published version we expand §5 dealing with relatives of the symmetric closure; note also that if K is an ordered field which is symmetrically closed or just has no cut of cofinality $(\text{cf}(K), \text{cf}(K))$ then K is real closed.

Our problem translates to considering cuts of K and their pair of cofinalities. Our strategy is:

Key words and phrases. Real closed fields, ordered fields, cuts, completeness, cofinality.

First typed: March 2000.

Research supported by the United States-Israel Binational Science Foundation. Publication 757.

- (a) we consider some properties of cuts (of a real closed field) being Dedekind, being Scott, being positive, being additive, being multiplicative,
- (b) we define dependence relation on the set of cuts of K , which satisfies the Steinitz assumptions,
- (c) realizing a maximal independent family of cuts with the right pairs of cofinality, we get a “one step symmetric closure”.

It is fine: we can show existence and uniqueness. But will iterating this “atomic closure” eventually terminate?

- (d) For a field K we define a similar chain inside K , its minimal length being $h(K)$
- (e) we define the depth $d(K)$ of K
- (f) we show that $h(K)$, and $d(K)$ are quite closed and show that our iterated closure from (c) do increase the cardinality much
- (g) we finally show that after $\leq \omega_2$ steps the iterated closure from (c) terminate.

2. REAL CLOSED FIELDS

Any ordered field embeds in a real closed field, and in fact has a unique real closure. We will find it convenient to work mainly with real closed fields throughout. Accordingly, we will need various properties of real closed fields. We assume some familiarity with quantifier elimination, real closure, and the like, and we use the following consequence of o -minimality. (Readers unfamiliar with o -minimality in general may simply remain in the context of real closed fields, or in geometrical language, *semialgebraic geometry*.)

Fact 2.1. (1) *Let K be a real closed field, and let f be a parametrically definable function of one variable defined over K . Then f is piecewise monotonic, with each piece either constant or strictly monotonic; that is we can find finitely many intervals (allowing ∞ and $-\infty$ as end points), f is constant strictly more or strictly decreasing on each interval. Moreover this holds uniformly and definably in definable families, with a bound on the number of pieces required, and with each piece an interval whose endpoints are definable from the defining parameters for the function.*

Notation 2.2. K, L are ordered fields usually real closed. If $K \subseteq L, X \subseteq L$, then $K(X)$ is the subfield of L generated by $K \cup X$. $K_+ = \{c \in K : c > 0\}$.

(2A) Cuts.

Definition 2.3. (1) A *cut* in a real closed field K is a pair $C = (C^-, C^+)$ with K the disjoint union of C^- and C^+ , and $C^- < C^+$. The cut is a *Dedekind cut* if both sides are nonempty, and C^- has no maximum, while C^+ has no minimum. For $L \subseteq K$ let $C|L = (C^- \cap L, C^+ \cap L)$, be a cut of L .

- (2) The *cofinality* of a cut C is the pair (κ, λ) with κ the cofinality of C^- and λ the coinitality of C^+ (i.e., the “cofinality to the left”). If the cut is not a Dedekind cut, then one includes 0 and 1 as possible values for these invariants.
- (3) A cut of cofinality (κ, λ) is *symmetric* if $\kappa = \lambda$.
- (4) A real closed field is *symmetrically complete* if it has no symmetric cuts.
- (5) A cut is *positive* if $C^- \cap K_+$ is nonempty.

Conclusion 2.4. (1) If L is a real closed field extending K and $a, b \in L \setminus K$ realizes the same cut of K then a, b realizes the same types over K in L .

- (2) If C is a non-Dedekind cut of K then for some θ , $\text{cf}(C) \in \{(\theta, 0), (0, \theta), (\theta, 1), (1, \theta)\}$ so $\theta = \text{cf}(K)$

Proof: By 2.1.

We will need to consider some more specialized properties of cuts.

Definition 2.5. Let K be a real closed field, C a cut in K .

- (1) The cut C is a *Scott cut* if it is a Dedekind cut, and for all $r > 0$ in K , there are elements $a \in C^-$, $b \in C^+$ with $b - a < r$.
- (2) The cut C is *additive* if C^- is closed under addition and contains some positive element.
- (3) The cut C is *multiplicative* if $C^- \cap K_+$ is closed under multiplication and contains 2.
- (4) C_{add} is the cut with left side $\{r \in K : r + C^- \subseteq C^-\}$.
- (5) C_{mlt} is the cut with left side $\{r \in K : r \cdot (C^- \cap K_+) \subseteq C^-\}$.

Observe that

Observation 2.6. (1) *Scott cuts are symmetric, in fact both cofinalities are equal to $\text{cf}(K)$.*

- (2) *If C is a positive Dedekind cut which is not a Scott cut, then C_{add} is an additive cut, or note: $C_{\text{add}}^- = \{c : c \leq 0\}$ is impossible as “not scott”*
- (3) *if C is an additive cut which is not a multiplicative cut, then C_{mlt} is a multiplicative cut.*

Definition 2.7. (1) If $K \subseteq L$ are ordered fields, then a cut C in K is said to be *realized*, or *filled*, by an element a of L if the cut induced by a on K is the cut C .

- (2) If $C_1, C_2 \subseteq K$ and $C_1 < C_2$ but no $a \in K$ satisfies $C_1 < a < C_2$ then the cut of K defined (or induce or canonically extends) by (C_1, C_2) is $(\{a \in K : a < c \text{ for some } c \in C_1\}, \{b \in K : c \leq b \text{ for some } c \in C_2\})$, e.g., (C_1, C_2) may be a cut of a subfield of K .

Lemma 2.8. *Let K be a real closed field. Then there is a real closed field L extending K in which every Scott cut has a unique realization, and no other Dedekind cuts are filled.*

This is called the *Scott completion* of K , and is strictly analogous to the classical Dedekind completion. The statement found in [?] is worded differently, without referring directly to cuts, though the relevant cuts are introduced in the course of the proof. The result is also given in greater generality there.

Lemma 2.9. *Let K be a real closed field, C a multiplicative cut in K , and L the real closure of $K(x)$, where x realizes the cut C . Then for any $y \in L$ realizing the same cut, we have $x^{1/n} < y < x^n$ for some n .*

Proof. Let $\mathcal{O}_K$ be $\{a \in K : |a| \in C^-\}$, and let $\mathcal{O}_L$ be the convex closure in L of $\mathcal{O}_K$. Then these are valuation rings, corresponding to valuations on K and L which will be called v_K and v_L respectively.

The value group Γ_K of v_K is a divisible ordered abelian group, and the value group of the restriction of v_L to $K(X)$ is $\Gamma_K \oplus \mathbb{Z}\gamma$ with $\gamma = v_L(x)$ negative, and infinitesimal relative to Γ_K . The value group of v_L is the divisible hull of $\Gamma_K \oplus \mathbb{Z}\gamma$.

Now if $y \in L$ induces the same cut C on K , then $v_L(y) = qv_L(x)$ for some positive rational q . Hence $u = y/x^q$ is a unit of $\mathcal{O}_L$, and thus $u, u^{-1} < x^\epsilon$ for all positive rational ϵ . So $x^{q-\epsilon} < y < x^{q+\epsilon}$ and the claim follows. $\square$

Lemma 2.10. *Let $K \subseteq L$ be real closed fields, and C an additive cut in L . Let C' and C'_{mlt} be the cuts induced on K by C and C_{mlt} respectively. Suppose that $C'_{\text{mlt}} = (C')_{\text{mlt}}$, and that $x, y \in L$ are two realizations of the cut C' , with $x \in C^-$ and $y \in C^+$. Then y/x induces the cut C'_{mlt} on K .*

Proof. If $a \in K$ and $ax \geq y$, then $a \in (C_{\text{mlt}})^+$, by definition, working in L .

On the other hand if $a \in K$ and $ax < y$, then $a \in [(C')_{\text{mlt}}]^-$, which by hypothesis is $(C'_{\text{mlt}})^-$. $\square$

Lemma 2.11. *Let $K \subseteq L$ be real closed fields, and C a positive Dedekind cut in L which is not additive. Let C' and C'_{add} be the cuts induced on K by C and C_{add} respectively. Suppose that $C'_{\text{add}} = (C')_{\text{add}}$. Suppose that $x, y \in L$ are two realizations of the cut C' , with $x \in C^-$ and $y \in C^+$. Then $y - x$ induces the cut C'_{add} on K .*

Proof. If $a \in K$ and $a + x \geq y$, then $a \in (C_{\text{add}})^+$, by definition, working in L .

On the other hand if $a \in K$ and $a + x < y$, then $a \in [(C')_{\text{add}}]^-$, which by hypothesis is $(C'_{\text{add}})^-$. $\square$

(2B) Independent cuts.

We will rely heavily on the following notion of independence.

Definition 2.12. Let K be a real closed field, and $\mathcal{C}$ a set of cuts in K . We say that the cuts in $\mathcal{C}$ are *dependent* if for every real closed field L containing realizations a_C ($C \in \mathcal{C}$) of the cuts over K , the set $\{a_C : C \in \mathcal{C}\}$ is algebraically dependent over K .

The following merely rephrases the definition.

Lemma 2.13. *Let K be a real closed field and $\mathcal{C}$ a set of cuts over K .*

- (1) *the following are equivalent.*
 - (A) *$\mathcal{C}$ is independent*
 - (B) *For each set $\mathcal{C}_0 \subseteq \mathcal{C}$, and each ordered field L containing K , if $a_C \in L$ is a realization of the cut C for each $C \in \mathcal{C}_0$, then the real closure of $K(a_C : C \in \mathcal{C}_0)$ does not realize any cuts in $\mathcal{C} \setminus \mathcal{C}_0$.*
- (2) *For $\mathcal{C}_0$ and L as in clause (B) above, every $C \in \mathcal{C} \setminus \mathcal{C}_0$ define a (unique) cut C' of L (see Definition 2.3(2)) and $\{C' : C \in \mathcal{C} \setminus \mathcal{C}_0\}$ is an independent set of cuts of L*
- (3) *Assume $\langle K_\alpha : \alpha \leq \delta \rangle$ is an increasing sequence of real closed fields, $\mathcal{C}$ a set of cuts of K_0 , and $C \in \mathcal{C} \wedge \alpha < \delta \Rightarrow C$ define a cut of K_δ . Then each $C \in \mathcal{C}$ define a cut of K_δ which we call $C^{[\kappa]}$ and if $\mathcal{C}$ is an independent set of cuts of K_0 then $\{C^{[\kappa]} : C \in \mathcal{C}\}$ is an independent set of cuts of K_δ .*

Note that this dependence relation satisfies the Steinitz axioms for a dependence relation. We will make use of it to realize certain sets of types in a controlled and canonical way.

Lemma 2.14. *Let K be a real closed field, and $\mathcal{C}$ a set of cuts over K . Then there is a real closed field L generated over K (as a real closed field) by a set of realizations of some independent family of cuts included in $\mathcal{C}$, in which all of the cuts $\mathcal{C}$ are realized. Furthermore, such an extension is unique up to isomorphism over K .*

Proof. Clearly we must take L to be the real closure of $K(a_C : C \in \mathcal{C}_0)$, where $\mathcal{C}_0$ is some maximal independent subset of $\mathcal{C}$; and equally clearly, this works.

It remains to check the uniqueness. This comes down to the following: for any real closed field L extending K , and for any choice of independent cuts $C_1, \dots, C_n$ in K which are realized by elements $a_1, \dots, a_n$ of L , the real closure of the field $K(a_1, \dots, a_n)$ is uniquely determined by the cuts. One proceeds by induction on n . The real closure $\hat{K}$ of $K(a_n)$ is determined by the cut C_n by 2.4; and as none of the other cuts are realized in it, they extend canonically to cuts $C'_1, \dots, C'_{n-1}$ over $\hat{K}$, which are independent over $\hat{K}$. At this point induction applies. $\square$

Lemma 2.15. *Let K be a real closed field, and $\mathcal{C}$ a set of Dedekind cuts in K . Suppose that C is a Dedekind cut of cofinality (κ, λ) which is dependent on $\mathcal{C}$, and let $\mathcal{C}_0$ be the set $\{C' \in \mathcal{C} : \text{cof}(C') = (\kappa, \lambda) \text{ or } (\lambda, \kappa)\}$. Then C is dependent on $\mathcal{C}_0$, and in particular $\mathcal{C}_0$ is nonempty.*

Proof. It is enough to prove this for the case that $\mathcal{C}$ is independent. If this fails, we may replace the base field K by the real closure $\hat{K}$ over K of a set of realizations of $\mathcal{C}_0$. Then since none of the cuts in $\mathcal{C} \setminus \mathcal{C}_0$ are realized, and

C is not realized, these cuts extend canonically to cuts over $\hat{K}$, and hence we may suppose $\mathcal{C}_0 = \emptyset$. We may also suppose $\mathcal{C}$ is finite, and after a second extension of K we may even assume that $\mathcal{C}$ consists of a single cut C_0 . This is the essential case.

So at this point we have a realization a of C_0 over the real closed field K , and a realization b of C over K , with b algebraic, and hence definable, over a , relative to K . Thus b is the value at a of a K -definable function (being the ℓ -th root for some i), not locally constant near a , and by Fact 2.1 it follows that there is an interval b with endpoints in K and a definable function which is order isomorphism or anti-isomorphism from the interval I_0 to an interval I_1 including a , with the cuts corresponding. So the cuts have some (or inverted) cofinalities. This contradicts the supposition that $\mathcal{C}_0$ has become empty, and proves the claim. $\square$

For our purposes, the following case is the main one. We combine our previous lemma with the uniqueness statement.

Proposition 2.16. *Let K be a real closed field, and $\mathcal{C}$ a maximal independent set of symmetric cuts in K . Let L be an ordered field containing K together with realizations a_C of each $C \in \mathcal{C}$. Then the real closure K' of $K(a_C : C \in \mathcal{C})$ realizes the symmetric cuts of K and no others. Furthermore, the result of this construction is unique up to isomorphism. Moreover if L' is a real closed field extending K which realizes every cut in $\mathcal{C}$ then K' can be embedded into L' over k .*

Proof: Clear (the “no other” by 2.18 below)

Evidently, this construction deserves a name.

- Definition 2.17.** (1) Let K be a real closed field. A *symmetric hull* of K is a real closed field generated over K by a set of realizations of a maximal independent set of symmetric cuts.
- (2) We say that $\bar{K} = \langle K_\alpha : \alpha \leq \alpha^* \rangle$ is an associated symmetric α^* -chain over K when:
- (a) $K_0 = K$,
 - (b) K_α is an order field ,
 - (c) K_α is increasing continuous with α
 - (d) $K_{\alpha+1}$ is a symmetric hull of K_α for $\alpha < \alpha^*$
- (3) In (2) we replace “ α^* -chain” by “chain” if K_{α^*} is symmetrically complete but $\alpha < \alpha^* \Rightarrow K_{\alpha+1} \neq K_\alpha$

While this is unique up to isomorphism, there is certainly no reason to expect it to be symmetrically complete, and the construction will need to be iterated. The considerations of the next section will help to bound the length of the iteration.

Lemma 2.18. (1) *For regular $\kappa < \lambda$ there is a real closed field with an (κ, λ) -cut*

- (2) Let K be a real closed field, and L its symmetric hull. Then every Scott cut in K has a unique realization in L .
- (3) Assume L is the real closure of $K \cup \{a_C : C \in \mathcal{C}\}$, $\mathcal{C}$ an independent set of cuts of K and a_C realizing the cut C in L for $C \in \mathcal{C}$.
 - (a) If every $C \in \mathcal{C}$ is a Dedekind cut then every cut of K realized in L is a Dedekind cut
 - (b) if one non-Dedekind cut of L is realized in L then every non-Dedekind cut of K is realized in L .

Proof. (1) First we choose K_i a real closed field increasing continuous with $i \leq \kappa$, $K_0 = K$ and for $i < \kappa$ the element $a_i \in K_{i+1} \setminus K_i$ is above all members of K_i . Second we choose a real closed field K^i increasing continuous with $i \leq \lambda$ such that $K^0 = K_\kappa$, and for $i < \lambda$, $b_i \in K_{i+1}^{i+1} \setminus K_i$ is above a_j for $j < \kappa$ and below any $b \in K^i$ such that $(\forall j < \kappa)(\ell_j < b)$, in K^i . Lastly in K^λ , $(\{a_j : j < \kappa\}, \{b_i : i < \lambda\})$ determine a cut i.e., $(\{a : a < a_j \text{ for some } j < \kappa\}, \{b : b_i < b \text{ for } i < \lambda\})$, in K_λ , is such a cut.

(2) Recall that every Scott cut is symmetric. One can form the symmetric hull of K by first taking its Scott completion K_1 , realizing only the Scott cuts (uniquely), and then taking the symmetric hull of K_1 , by part (3) we are done.

(3) Easy. □

3. HEIGHT AND DEPTH

Definition 3.1. Let K be a real closed field.

- (1) The *height* of K , $h(K)$, is the least ordinal α for which we can find a continuous increasing sequence K_i ($i \leq \alpha$) of real closed fields with K_0 countable, $K_\alpha = K$, and K_{i+1} generated over K_i , as a real closed field, by a set of realizations of cuts which is independent. [[we could have chosen K_0 as the algebraic members of K]]
- (2) Let $h^+(K)$ be $\max(|h(K)|^+, \aleph_1)$

Remark:

- (1) $\aleph_1$ is the first uncountable cardinal
- (2) $h^+(K)$ is the first uncountable cardinal strictly greater than $h(K)$, so regular.

Observe that the height of K is an ordinal of cardinality at most $|K|$ (or is undefined, you can let it be ∞ , a case which by 3.3 do not occurs). We need to understand the relationship of the height of K with its order-theoretic structure, which for our purposes is controlled by the following parameter.

Definition 3.2. Let K be a real closed field. The *depth* of K , denoted $d(K)$, is the least regular cardinal κ greater than the length of every strictly increasing sequence in K .

Observe that the depth is uncountable. The following estimate is straightforward, and what we will really need is the estimate in the other direction, which will be given momentarily.

Lemma 3.3. *Let K be a real closed field. Then $h(K) \leq d(K)$.*

Proof. One builds a continuous strictly increasing tower K_α of real closed subfields of K starting with any countable subfield of K . If α is limit, we define $K_\alpha = \bigcup_{\beta < \alpha} K_\beta$. For successor ordinals, $K_{\alpha+1}$ is the real closure of $K_\alpha \cup \{a_C^\alpha : C \in \bar{\mathcal{C}}_\alpha\}$ inside K where $\mathcal{C}_\alpha$ is a maximal set of symmetric cuts of K_α realized in K , and $a_C^\alpha \in K$ realized C . We stop when $K_\alpha = K$. If this continues past $\kappa = d(K)$, then there is a cut over K_κ filled at stage κ by an element $x \in K$. Then the cut determined by x over each K_α for $\alpha < \kappa$ is filled at stage $\alpha + 1$ by an element y_α . Those y_α lying below x form an increasing sequence, by construction, which is therefore of length less than κ ; and similarly there are fewer than κ elements $y_\alpha > x$, so we arrive at a contradiction. $\square$

Proposition 3.4. *Let K be a real closed field. Then $d(K) \leq h^+(K)$.*

Proof. Let $\kappa > h(K)$ be regular and uncountable, and let K_α ($\alpha \leq h(K)$) be a continuous increasing chain of real closed fields, with K_0 countable, $K_{h(K)} = K$, and K_{i+1} generated over K_i , as a real closed field, by a set of realizations of independent cuts.

For $\alpha \leq h(K)$ and $X \subseteq K$, let $K_{\alpha,X}$ be the real closure of $K_\alpha(X)$ inside K . We recast our claim as follows to allow an inductive argument.

⊗ For $X \subseteq K$ with $|X| < \kappa$, and any $\alpha \leq h(K)$, we have $d(K_{\alpha,X}) \leq \kappa$.

Now this claim is trivial for $\alpha = 0$ as K_0 is countable, and the claim passes smoothly through limit ordinals up to $h(K)$, so we need only consider the passage from α to $\beta = \alpha + 1$. So K_β is $K_{\alpha,S}$ with S a set of realizations of independent cuts over K_α , and similarly $K_{\beta,X}$ is $K_{\alpha,X \cup S}$.

Consider the claim in the following form:

$$d(K_{\alpha,X \cup S_0}) \leq \kappa \text{ for } S_0 \subseteq S$$

In this form, it is clear if $|S_0| < \kappa$, as it is included in the inductive hypothesis for α , and the case $|S_0| \geq \kappa$ reduces at once to the case $|S_0| = \kappa$. So we now assume that S_0 is a set of realizations of independent cuts of K_α .

By 2.13(2),(3) we can find a subset S_1 of S_0 of cardinality $\aleph_0 + |X|$ such that:

- (a) if $s \in S_0 \setminus S_1$ then the cut C which s induce on K_α is not realized in the real closure K'_α of $K_\alpha(X_0 \cup S_1)$
- (b) the cuts which the $s \in S_0 \setminus S_1$ induce on K'_α form an independent family then after moving S_1 into X , we may suppose that S_0 is a set of realizations of cuts which are independent over $K_{\alpha,X}$.

Let $\{s_\epsilon : \epsilon < \kappa\}$ list $S_0 \setminus S_1$. For $\zeta \leq \kappa$, let $L_\zeta = K_{\alpha, X \cup \{s_\epsilon : \epsilon < \zeta\}}$ and let $L = L_\kappa$. By the induction hypothesis (for α and $X \cup \{s_\epsilon : \epsilon < \zeta\}$) we have $d(L_\zeta) \leq \kappa$ for $\zeta < \kappa$, and we shall prove $d(L) \leq \kappa$.

Let C_i be the cut realized by s_i over L_0 . Note that C_i extends canonically to a cut C_i^j on K_j for all $j \leq i$, and for fixed j , the set $\{C_i^j : i \in [j, \kappa)\}$ of cuts is independent.

Now suppose, toward a contradiction, that we have $(a_i : i < \kappa)$ increasing in L . Necessary for each $i < \kappa$ for some ordinal $f(i) < \kappa$, $a \in L_{f(i)}$, so $f : \kappa \rightarrow \kappa$ is well defined and for each $j < \kappa$ the set $\{i < \kappa : a_1 \in L_j\}$ is a bounded subset of κ . Without loss of generality f is increasing. Now for i with $f(i) > \epsilon$ let B_i^ϵ denote the cut induced on L_ϵ by a_i . With ϵ held fixed, and with i varying, as $d(L_\epsilon) \leq \kappa$ we find that the cuts B_i^ϵ stabilize for large enough $i < \kappa$ (and furthermore, $a_i \notin L_\epsilon$). Accordingly, for each ϵ we may select $j_\epsilon < \kappa$ above j_ϵ^0 such that the cuts B_i^ϵ coincide for all $i \geq j_\epsilon$.

Now fix a limit ordinal $\delta < \kappa$ such that for all $\epsilon < \delta$ we have $j_\epsilon < \delta$. We may also require that $a_i \in L_\delta$ for $i < \delta$. Then $(B_\delta^\delta)^- = \bigcup_{\epsilon < \delta} (B_{j_\epsilon}^\epsilon)^-$, and the cofinality from the left of B_δ^δ is $\text{cof}(\delta)$.

Now a_δ is algebraic over $L_\delta(s_i : i \in I_0)$ for some finite subset I_0 of $[\delta, \kappa)$, and hence also over $L_\epsilon(s_i : i \in I_0)$ for some $\epsilon < \delta$. Thus the cut B_δ^ϵ depends on the cuts C_i^ϵ ($i \in I_0$) over L_ϵ . As $j_\epsilon < \delta$ necessarily $B_\delta^\epsilon = B_{j_\epsilon}^\epsilon$ is realized in L_δ , it follows that this cut is also dependent on the sets $\{C_i^\epsilon : i < \delta\}$ of cuts over L_ϵ . But the cuts C_i^ϵ for $i \geq \epsilon$ are supposed to be independent over L_ϵ , a contradiction. $\square$

Proposition 3.5. *Let K be a real closed field. Then $|h(K)| \leq |K| \leq 2^{|h(K)|}$.*

Proof. The first inequality is clear. For the second, let $\alpha = h(K)$, $\kappa = |\alpha| + \aleph_0$, and let K_i ($i \leq \alpha$) be a chain of the sort afforded by the definition of the height. Note that $h^+(K) = \kappa^+$. We show by induction on i that $|K_i| \leq 2^\kappa$. Only successor ordinals $i = j + 1$ require consideration, where we suppose $|K_j| \leq 2^\kappa$.

Each generator a of K_i over K_j corresponds to a cut C_a in K_j , and each such cut is determined by the choice of some cofinal sequence S_a in C_a^- . Such a sequence S_a may be taken to have order type a regular cardinal, and will have length less than $d(K)$. Since $d(K) \leq h^+(K)$, we find that the order type of S_a is at most κ . So the number of such sequences is at most $\sum_{\lambda \leq \kappa} |K_j|^\lambda \leq \kappa \times (2^\kappa)^\kappa = 2^\kappa$. $\square$

Claim 3.6. *By [1], in our terms.*

- (1) *If L is the Scott completion of K (i.e., in our terms as in definition 2.5(1). But by $\mathcal{C}$ is a maximal set of Scott cuts of K which is independent, then K*
 - (a) *K is dense in L*
 - (b) *for every Dedekind cut C of L the cut $C \restriction K$ induce C hence $\text{cf}(C) = \text{cf}(C \restriction K)$*

- (2) If K has no symmetric cut except possibly Scott cuts, and L is a symmetric closure of K (equivalently the Scott completion of K) then L is symmetrically complete.

Proof: Striaight.

4. PROOF OF THE THEOREM

We now consider the following construction. Given a real closed field K , we form a continuous increasing chain K_α by setting $K_0 = K$, taking $K_{\alpha+1}$ to be the symmetric hull of K_α in the sense of Definition 2.17, and taking unions at limit ordinals.

If at some stage K_α is symmetrically complete, that is $K_\alpha = K_{\alpha+1}$, then we have the desired symmetrically complete extension of K , and furthermore our extension is prime in a natural sense. We claim in fact:

- Proposition 4.1.** (1) Assume K is a real closed field, $\kappa = \aleph_2, \kappa' = \max(h^+(K), \aleph_2)$ and K_α ($\alpha \leq \kappa + 1$) is an associated continuous symmetric κ -chain, then
- (a) if $\text{cf}(K) \neq \kappa$ then K_κ is symmetrically complete so $K_{\kappa+1} = K_\kappa$.
 - (b) $K_{\kappa+1}$ is symmetrically closed
- (2) Also
- (i) $|K_\kappa| \leq 2^{h^+(K) + \aleph_1}$, and $|K_{\kappa+1}| \leq 2^{h^+(K) + \aleph_1}$
 - (ii) if K' is a symmetrically complete extension of K then $K_{\kappa+1}$ can be embedded into K' over K .
 - (iii) K is unbounded in $K_{\kappa+1}$ (and no non-Dedekind cut of K is realized in K_κ and no non-symmetric Dedekind cut of K is realized in K_κ).
 - (iv) Any two real closed fields extending K which are symmetrically complete and embeddable into each other, are isomorphism over K (so we can say K_κ is the closure)
 - (v) For some unique $\alpha^* \leq \kappa$ there is an associated continuous chain over K of length α^* .

The proof of Proposition 4.1 occupies the remainder of this section.

Lemma 4.2. Suppose that K is a real closed field, and that $\langle K_\alpha : \alpha \leq \alpha(*) \rangle$ is a continuous chain of iterated symmetric hulls of any length. Let $x \in K_\alpha \setminus K$ with $\alpha > 0$ arbitrary. Then the cut induced on K by x is symmetric.

Proof. Let β be minimal such that the cut in question is filled in $K_{\beta+1}$. Then the cut induced on K_β by x is the canonical extension of the cut induced on K by x , and is symmetric by Proposition 2.16. $\square$

We now begin the proof by contradiction of Proposition 4.1(1). We assume therefore that the chain is strictly increasing at every step up to K_κ , and that there is a symmetric cut C over K_κ .

Remark: If we use $\kappa = \max\{h^+(K), \aleph_2\}$ then $\kappa \geq \aleph_2$ is regular and greater than $h(K)$; in particular $\kappa \geq d(K)$ by 3.4. Furthermore, as $\kappa > h(K)$, we can view the chain K_α as a continuation of a chain $\hat{K}_i$ ($i \leq h(K)$) of the sort occurring in the definition of $h(K)$, with $\hat{K}_{h(K)} = K_0$; then the concatenated chain gives a construction of K_α of length at most $h(K) + \alpha < \kappa$, and hence $h(K_\alpha) < \kappa$ for all $\alpha < \kappa$, and in particular $d(K_\alpha) \leq \kappa$ for all $\alpha < \kappa$ by 3.4.

For $\alpha < \kappa$, let C_α denote the cut induced on K_α by C .

Lemma 4.3. (1) *For any $\alpha < \kappa$, the cut C_α is symmetric.*
 (2) *For every $\alpha < \kappa$, C_α^- is bounded in C^- and C_α^+ is bounded in C^+ from below.*

Proof:

- (1) Suppose C_α is not symmetric. Then the cut C_α is not realized in K_κ , by Lemma 4.2. Hence the cut C is the canonical extension of C_α to K_κ , contradicting its supposed symmetry.
- (2) Assume C_α^- is unbounded from below in C^- ; so necessarily $\text{cf}(C_\alpha^-) \equiv \kappa$ and $\beta \in [\alpha, \kappa) \Rightarrow \text{cf}(C_\beta^-) = \text{cf}(C_\alpha^-)$.

For every $\beta \in [\alpha, \kappa)$, as $K_{\alpha+1}$ is a symmetric hule of K , by part (1) it follows that some $a_\beta \in K_{\alpha+1}$ realizes the cut C_β hence by the previous sentence, $a_\beta \in C^+$. So for every limit $\delta < \kappa$, $\delta \in E \Rightarrow \{a_\beta : \beta \in [\alpha, \delta)\}$ is a subset of C_δ^+ unbounded from below, hence $\text{cf}(C_\delta) = (\text{cf}(C_\delta^-), \text{cf}(C_\delta^+)) = (\text{cf}(C_\alpha^-), \text{cf}(\delta))$. As $\aleph_0 \neq \aleph_1$ are regulars $< \kappa$ for some $\delta \in (\alpha, \kappa)$ $\text{cf}(C_\alpha^-) \neq \text{cf}(\delta)$ hence C_δ is not symmetric, contradicting part (1).

In particular, the cut C_α is realized in $K_{\alpha+1}$, and thus we have the following.

Corollary 4.4. *For any limit ordinal $\alpha \leq \kappa$, the two-sided cofinality of C_α is $\text{cof}(\alpha)$.*

After these preliminaries, we divide the analysis of the supposed cut C into a number of cases, each of which leads to a contradiction. Let $\langle C_\zeta^* : \zeta < \theta \rangle$ be an increasing sequence of members of K_+ cofinal in it.

(Case I) C is a Scott cut

If C is a Scott cut and $\text{cf}(K) = \aleph_2$, there is nothing to be proved, so assume $\text{cf}(K) \neq \aleph_2$.

By 4.3(2), we can find $\langle a_\alpha^-, a_\alpha^+ : \alpha < \kappa \rangle$, such that $a_\alpha^- \in C^-$, $a_\alpha^+ \in C^+$ both realizes the cut C_α . For some club E of κ consisting of limit ordinals we have $\alpha < \delta \in E \Rightarrow a_\alpha^-, a_\alpha^+ \in K_\delta$. As C is a Scott cut necessarily $\langle a_\alpha^- - a_\alpha^+ : \alpha < \kappa \rangle$ is a decreasing sequence of negative members of K_κ with no negative lower bound, so $\langle 1/(a_\alpha^+ - a_\alpha^-) : \alpha < \kappa \rangle$ is increasing cofinal in K_κ , so $\text{cf}(K_\kappa) = \kappa$. But K is cofinal in K_κ hence $\text{cf}(K) = \kappa = \aleph_2$, contradicting what we have assumed in the beginning of the case.

(Case II) C is a multiplicative cut

Let $\alpha < \kappa$ have uncountable cofinality (recall $\kappa \geq \aleph_2$).

The cut C_α is realized in $K_{\alpha+1}$ by some element a . As C is multiplicative, either all positive rational powers of a lie in C^- , or all positive rational powers of a lie in C^+ .

On the other hand, $K_{\alpha+1}$ may be constructed in two stages as follows. First realize all the cuts in a maximal independent set of symmetric cuts in K_α , with the exception of the cut C_α , getting a field K'_α ; then take the real closure of $K'_\alpha(a)$, where a fills the canonical extension of the cut C_α to K'_α . As seen in Lemma 2.9, there are only two cuts which may possibly be induced by C on $K_{\alpha+1}$, and each has countable cofinality from one side, and uncountable cofinality from the other.

So $C_{\alpha+1}$ is not symmetric, and this is a contradiction.

(Case III) C is an additive cut

Let $\langle b_\alpha^- : \alpha < \kappa \rangle$ be increasing cofinal in $C^- \cap K_+$ and let $\langle b_\alpha^+ : \alpha < \kappa \rangle$ be decreasing unbounded from below in C^+ . Clearly $\alpha < \kappa \Rightarrow b_\alpha^+ / b_\alpha^- \in (C_{\text{mlt}})^+$ and easily $\langle b_\alpha^+ / b_\alpha^- : \alpha < \kappa \rangle$ is a decreasing sequence of members of $(C_{\text{mlt}})^+$ unbounded from below in it

(Quotation ??)

For some club E of κ we have: if $\delta \in E$ then $\langle b_\alpha^- : \alpha < \delta \rangle$ is increasing sequence in $C^- \cap (K_\delta)_+$ cofinal in it and $\langle b_\alpha^+ : \alpha < \delta \rangle$ is a decreasing sequence in $C^+ \cap K_\delta$ unbounded from below in it. So $\langle b_\alpha^+ / b_\alpha^- : \alpha < \delta \rangle$ is a decreasing sequence of members of $(C_{\text{mlt}})^+$ unbounded from below in it. [[Also it is a decreasing sequence of members of $(C \upharpoonright K_\delta)_{\text{mlt}}^+$. Hence $(C_{\text{mlt}})^+ \cap K_\delta = (C \upharpoonright K_\delta)_{\text{mlt}}^+$ i.e., $(C_\delta)_{\text{mlt}} = (C_{\text{mlt}})_\delta$, recalling 2.5(5).]]

According to the property of $\langle b_\alpha^+ / b_\alpha^- : \alpha < \kappa \rangle$ the cofinality of (C_{mlt}) from the right is κ .

Now if the cofinality of C_{mlt} from the left is also κ , then by 2.6(3) we contradict Case II. On the other hand if the cofinality of C_{mlt} from the left is less than κ , then from some point downward this cofinality stabilizes; but then, choosing δ large and of some other cofinality (again, since $\kappa \geq \aleph_2$ this is possible), we contradict Lemma 4.3.

(Case IV) C is a positive Dedekind cut, but not a Scott cut

One argues as in the preceding case, considering C_{add} and using Lemma 2.11, which leads to a symmetric additive cut and thus a contradiction to the previous case. In details choose $\langle b_\alpha^- : \alpha < \kappa \rangle, \langle b_\alpha^+ : \alpha < \kappa \rangle$ as in case III, so $\langle b_\alpha^+ - b_\alpha^- : \alpha < \kappa \rangle$ is a decreasing sequence in C_{add}^+ unbounded from below in it

(Quotation ?)

[[As in (case III) for a club of $\delta < \kappa$, $(C_{\text{add}})^+ \cap K_\delta = (C \upharpoonright K_\delta)_{\text{add}}^+$ hence $(C_\delta)_{\text{mlt}} = (C_{\text{mlt}})$]]. If the cofinality of C_{add} from below is also κ , recall that C_{add} is an additive cut by 2.6(2) contradiction to case III.

As no cases remain, Proposition 4.1(1) is proved, and thus the construction of a symmetrically complete extension terminates.

As for clause (i) of 4.1(2), to estimate the cardinality of the resulting symmetrically complete extension, recall that it has height at most $\kappa' = \max(h^+(K), \aleph_2) \leq \max(|K|^+, \aleph_2)$ and hence cardinality at most $2^{\kappa'}$. Moreover, similarly for any $\alpha < \kappa'$, $|K_\alpha| \leq 2^{h^+(K) + \aleph_1}$ hence $|K| = |\bigcup_{\alpha < \kappa} K_\alpha| \leq \sum_{\alpha < \kappa} |K_\alpha| \leq \sum_{\alpha < \kappa} 2^{h^+(K) + \aleph_1} = \kappa' + 2^{h^+(K) + \aleph_1} = 2^{h^+(K) + \aleph_1}$

For clause (ii) of 4.1, we define an embedding h_α of K_α into K' , increasing continuous with α for $\alpha \leq \kappa$. For $\alpha = 0$, h_0 is the identity, for α limit take union and for $\alpha = \beta + 1$ use 2.16.

Clauses (iii),(iv) of 4.1 is easy too.

5. CONCLUDING REMARKS

Discussion: It should be clear that there are considerably more general types of closure that can be constructed in a similar manner. Let Θ be a class of possible cofinalities of cuts, that is pairs of regular cardinals, and suppose that Θ is symmetric in the sense that $(\theta_1, \theta_2) \in \Theta$ implies $(\theta_2, \theta_1) \in \Theta$. Then we may consider Θ -constructions in which maximal independent sets of cuts, all of whose cofinalities are restricted to lie in Θ , are taken. In order to get such a construction to terminate, all that is needed is the following:

- (a) for all regular θ_1 , there is θ_2 such that the pair (θ_1, θ_2) is not in Θ .
- (b) for some regular $\kappa \geq h(K) + \aleph_2$, for every θ_1 regular $< \kappa$ there is a $\theta_2 < \kappa$ such that $(\theta_1, \theta_2) \notin \Theta$

The proof is as above; in the symmetric case, Θ_{sym} consists of all pairs (θ, θ) of equal regular cardinals. Clearly we may make the closure to be equal large as we need κ as in (4) above. Also in the proof of 4.1(1) in the multiplicative case we choose δ such that $(\aleph_0, \text{cf}(\delta)) \notin \Theta$ (but of course change the cardinality bound).

Under the preceding mild conditions, such a Θ -construction provides an “atomic” extension of the desired type. So we have Θ -closure, and it is prime (as in clause (ii) of 4.1(2)). We also can change the cofinality of K .

Question:

- (1) $h(K)$ unique: ?? ordered field $(\text{cf} K, \text{cf} K) \notin \Theta$, not real closed.
- (2) See “Vive la difference II” complete Scott

Definition 5.1. Assume Θ is a set or class of pairs of regular cardinals which is symmetric i.e., $(\kappa, \lambda) \in \Theta \Rightarrow (\lambda, \kappa) \in \Theta$.

- (0) A Θ -cut of K is a cut C with $\text{cf}(C) \in \Theta$

- (1) We say that a real closed field is Θ -complete if no Dedekind cut of K has cofinality $\in \Theta$
- (2) We say that L is a Θ -hull of K when there is a maximal subset $\mathcal{C}$ of $\text{cut}_\Theta(K) := \{C : C \text{ a Dedekind cut of } K \text{ with cofinality } \in \Theta\}$ which is independent and $a_C \in L$ for $C \in \mathcal{C}$ realizing C such that L is the real closure of $K \cup \{a_C : C \in \mathcal{C}\}$
- (2A) We say that L is a weak Θ -hull of K when there is a subset $\mathcal{C}$ of $\text{cut}_\Theta(K)$ which is independent and $a_C \in L$ for $C \in \mathcal{C}$ realizing C such that L is the real closure of $K \cup \{a_C : C \in \mathcal{C}\}$
- (3) We say that $\langle K_\alpha : \alpha \leq \alpha^* \rangle$ is an associated $\Theta - \alpha^*$ -chain over K when: K_α is a real closed field, increasing continuous with α , $K_0 = K$ and $K_{\alpha+1}$ is a Θ -hull of K_α for $\alpha < \alpha^*$
- (3A) We say that $\langle K_\alpha : \alpha \leq \alpha^* \rangle$ is a weak associated $\Theta - \alpha^*$ -chain over K when: K_α is increasing continuous sequence of real closed fields, $K = K_0$ and $K_{\alpha+1}$ a weak Θ -hull of K_α for $\alpha < \alpha^*$. We may omit Θ meaning $\{(\kappa, \lambda) : \kappa, \lambda \text{ are regular infinite cardinals}\}$. We may omit “over K ”
- (4) We say that $\langle K_\alpha : \alpha \leq \alpha^* \rangle$ is an associated Θ -chain over K when: it is an associated $\Theta - \alpha^*$ -chain over K , $\alpha < \alpha^* \Rightarrow K_{\alpha+1} \neq K_\alpha$ and K_{α^*} is Θ complete
- (5) Let $d'(K)$ be the minimal regular cardinal κ (so infinite such that: for every non-Scott Dedekind cut C of K both cofinalities of C are $< \kappa$).

Theorem 5.2. *Let K be a real closed field and Θ be as in Definition 5.1.*

- (1) *There is a Θ -hull L of K , see Definition 5.1(1)*
- (2) *L in (1) is unique up to isomorphism over K , and K is cofinal in it*
- (3) *For every ordinal α^* there is an associated $\Theta - \alpha^*$ -chain over K , see Definition 5.1(2) and it is unique up to isomorphism over K*
- (3A) *If $\langle K_\alpha : \alpha \leq \alpha^* \rangle$ is a weak associated $\Theta - \alpha^*$ -chain over this K it is cofinal in all its members in particular in K_{α^*}*
- (4) *If $\theta = \text{cf}(\theta) < d(K)$ moreover for some λ , K has a non-Scott Dedekind cut of cofinality (θ, λ) and $(\forall \lambda)[(\theta, \lambda) \in \Theta]$ then there is no associated Θ -chain over K , moreover no Θ -complete extension of K*
- (5) $\aleph_0 \leq d'(K) \leq d(K)$
- (6) *If $d'(K) = \aleph_0$ then K is isomorphic to the field of the reals*
- (7) *If $\langle K_\alpha : \alpha \leq \alpha^* \rangle$ is a weak associated $\Theta - \alpha^*$ -chain over K then $d'(K) \leq \text{Max}\{|\alpha^*|^+, d'(K)\}$,
Check*
- (7A) *In (7) also $|K_{\alpha^*}| \leq |K| + |\alpha^*| + \sum\{|K|^\theta : (\theta, \theta) \in \Theta \text{ and } \Theta < d'(K)\}$*
- (8) *Assume κ satisfies “ $\kappa \neq \text{cf}(K)$, $\kappa = \text{cf}(\kappa) > \aleph_0$ and for every regular $\theta < d'(K) + \kappa$ for some regular $\lambda < \kappa$ we have $(\theta, \lambda) \notin \Theta$, then there is an associated Θ -chain over K of length $\leq \kappa$ (compare with 4)*
- (9) *If $(\forall \text{ regular } \theta) (\exists \text{ regular } \lambda) [(\theta, \lambda) \notin \Theta]$ then for every real closed field K' there is an associated Θ -chain over it*

- (10) *If there is an associated Θ -chain $\langle K_\alpha : \alpha \leq \alpha^* \rangle$ over K then*
- (a) *K_{α^*} is a real closed field, Θ -complete extending K*
 - (b) *[universally] if K' is a real closed field extending K then K_{α^*} can be embedded into K' over K*
 - (c) *[uniqueness] if L is a Θ -complete real closed field extending K which can be embedded over K into L' for every Θ -complete, Θ -complete real closed field extending K then L is isomorphism to K_{α^*} over K .*
 - (d) *in (c), $|L| \leq 2^{<\kappa}$, κ minimal as in (8).*

Proof:

- (1) As in the proof of xxx
- (2) As in the proof of yyy
- (3) Follows from (1)+(2)
- (3A) Easily by induction on α^* .
- (4) Let C be a non-Scott cut of K such that $\text{cf}(C) = (\theta, \lambda)$, without loss of generality C is positive and $\langle a_\alpha : \alpha < \theta \rangle$ be an increasing sequence of members of $C \cap K_+$ cofinal in it. As C is non-Scott clearly for some $c \in K_+$ we have $a \in C^- \Rightarrow a + c \in C^-$ hence without loss of generality

(*) $c \in K_+$ and $a_\alpha + c < a_{\alpha+1}$ for $\alpha < \theta$.

Now if L is a Θ -complete extension of K then there is a cut C_1 of L such that $C_1^- = \{a \in L : a < a_\alpha \text{ for some } \alpha < \theta\}$. Let $\text{cf}(C_1) = (\lambda_1, \lambda_2)$, so necessarily $\lambda_1 = \theta$ and obviously C_1^+ cannot have a first element b as then $b - c \in C_1^+$ by (*), so $\lambda_2 \geq \aleph_0$. So by an assumption

$(\lambda_1, \lambda_2) \in \Theta$, contradiction to “ L is Θ – complete”

- (5) Trivial, see Definition 5.1(xxx).
- (6) Easily K is complete hence it is well known to be isomorphism to the field of reals.
- (7) So (*)
 - (a) $\langle K_\alpha : \alpha \leq \alpha^* \rangle$ is a weak Θ -sequence
 - (b) K_0 is unbounded in K_{α^*} . Now we repeat the proof of 3.4

Older alternative

We prove by induction on $\alpha \leq \alpha^*$ that $(*)_\alpha$ for any Dedekind cut C of K_α of cofinality (θ, λ) we have $\theta + \lambda < d'(K) + |\alpha|^+$. The case $\alpha = 0$ holds by an assumption and the case α limit holds. So let $\alpha = \beta + 1$ and let $K_{\alpha+1}$ be the real closure of $K \cup \{a_C : C \in \mathcal{C}\}$ which is an independent family of Θ -cuts of K and downward continuous assume that $C, (\theta, \lambda)$ form a counter example so without loss of generality $\lambda \geq d'(k) + |\alpha|^+$

- (7A) By ?? quoted below, $\text{cut}_{\{(\theta, \lambda) : \theta \neq \lambda_{\text{regular}}\}}(K)$ has cardinality $\leq |K|$.
- (8) $(*)_1$ like 4.3 replacing “symmetric” cut by “ Θ -cut”. Let C be a cut of κ which is a Θ -cut with ??. If for some $\alpha < \kappa$ $C \upharpoonright K_\alpha$

induce C on K_κ , then $C \restriction K_\alpha$ is a cut of K_α of cofinality the same as C , hence a Θ -cut of K_α hence is realized in $K_{\alpha+1}$ by the construction, say by a , contradiction to “ $C \restriction K_\alpha$ induce C on K_κ ”. Hence

(*)₂ for no $\alpha < \kappa$ does $C \restriction K_\alpha$ induce C on K_κ . This means that for every $\alpha < \kappa$ some $a_\alpha \in K \setminus K_\alpha$ realizes C , so $a_\alpha \in C^- \vee a_\alpha \in C^+$ so as we can replace C by $|\{-b : b \in C^+\}, \{-b : b \in C^-\}|$ without loss of generality for arbitrarily large $\alpha < \kappa, a_\alpha \in C^-$, so remain ??

(*)₃ $a_\alpha \in C^- \subseteq K_\kappa$ realizes the cut $C \restriction K_\alpha$

(*)₄ $\text{cf}(C^-) = \kappa$ so let $\text{cf}(C) = (\kappa, \lambda)$

Case 1: If $\lambda \neq \kappa$ let σ be regular $< \kappa$ such that $(\sigma, \lambda) \notin \Theta$. Let $\{b_\beta : \beta < \lambda\}$ be a decreasing sequence in C^+ unbounded from below in it so without loss of generality

(*) $\{b_\beta : \beta < \lambda\} \subseteq k_{\alpha(*)}$

Now for some club E of κ we have

(*) if $\delta \in E$ then $\delta > \alpha(*)$ and $\{a_\alpha : \alpha < \delta\}$ is an unbounded subset of $C^- \cap K_\delta$ hence $\text{cf}(C \restriction K_\delta) = (\text{cf}(\delta), \lambda)$

Choose $\delta \in E$, such that $\text{cf}(\delta) = \sigma$. But then a_δ realizes a $\{(\text{cf}(\delta), \lambda)\}$ -cut i.e., $\{(\sigma, \lambda)\}$ -cut i.e., $\{(\sigma, \lambda)\}$ -cut which is a non Θ -cut by the choice of σ , contradiction to (*)₁.

Case 2: $\lambda = \kappa$

We repeat the proof of 4.1 after 4.3, in (case I) using $\kappa \neq \text{cf}(K)$

(9) For every field K' define κ_n by induction on $n < \omega'$

$$\kappa_0 = |K'| \text{ (or } d'(K))$$

$\kappa_n = \text{Min}\{\kappa \text{ regular and if } \theta < \kappa_n \text{ then for some } \lambda < \kappa_{n+1} \text{ we have } (\theta, \lambda) \notin \Theta\}$.

Now $K', (\sum\{\kappa_n : n < \omega\})^+$ satisfies the condition in (8).

(10) Easy. [Detials?]

What about $\text{cf}(K)$, we have not changed it in all our completion. It doesn't make much difference as ??.

Claim 5.3. (1) For K and regular $\kappa (\geq \aleph_0)$ there is L such that

(a) $K \subseteq L$

(b) $\text{cf}(L) = \kappa$

(c) if $K \subseteq L'$ and $\text{cf}(L') = \kappa$ then we can embed L into L' over K

(d) if $\kappa > \aleph_0$ then in clause (c) we can add: there is an embedding f of L into L' over K such that $\text{Rang}(f)$ is unbounded in L'

(2) For K and $\kappa = \aleph_0$ there are L_1, L , such that:

(a) $K \subseteq L_\ell$

(b) $\text{cf}(L) = \aleph_0$

(c) $\text{cf} K \subseteq L'$ then for some $\ell \in \{1, 2\}$ there is an embedding f of L into L' over K such that $\text{Range}(\ell)$ is unbounded in L'

(3) If $\text{cf}(K) > \aleph_0$ then in part (2) we cannot have $L_1 = L_2$,

- (4) If $\text{cf}(K) = \aleph_0$ then in part (2) we can use $L_1 = K = L_2$ similarly in part (1).

* * *

We have concentrated on real closed fields. This is justified by

- Claim 5.4.** (1) Assume K is an ordered field, $\theta = \text{cf}(K)$ and K has no $\{(\theta, \theta)\}$ -cut. Then K is real closed.
 (2) In Theorem 5.2 if we add $(\text{cf}(K), \text{cf}(K)) \notin \Theta$ and deal with ordered fields, it still holds.

- Claim 5.5.** (1) If F is an ordered field of cardinality μ then there is F' such that
 (a) F' is a real closed field F' of cardinality λ
 (b) F' extends F
 (c) $\text{cf}(F) = \aleph_0$
 (d) if C is a Dedekind cut of F' of cofinality (θ_1, θ_2) then $\theta_1 = \theta_2$
 (2) If I is a linear order of cardinality μ then we can find F such that
 (a) F is a real closed field of cardinality μ
 (b) I is embeddable into F (and if it is $(F_0^+, <)$, F' an ordered field we can add that F_0^+ is embeddable into
 (c) $\text{cf}(F) = \aleph_0$
 (d) if C is a Dedekind cut of F cofinality (θ_1, θ_2) then $\theta_1 = \theta_2$
 (e) Assume $\mu < \lambda$, and for every $J_1 \in [I]^{\leq \mu}$.

Then

- (α) $\mathcal{E}_\mu$ has $\leq 2^\mu$ e.q. there is J_2 , such that $J_1 \subseteq J_2 \in [I]^{\leq \mu}$ and every cut of I classes where $\mathcal{E}_\mu\{(\bar{a}, \bar{b})$ induce by a member of $I \setminus J_2$ has both cofinalities $\leq \aleph_0 \bar{a}, \bar{b} \in {}^\mu F$ and there is an order-auto of F mapping $\bar{a}$ to $\bar{b}$
 (β) (as in [1] fill).

REFERENCES

- [1] Saharon Shelah. Categoricity and solvability of A.E.C., quite highly. *preprint*. 0808.3023.
 [2] Saharon Shelah. Vive la différence II. The Ax-Kochen isomorphism theorem. *Israel Journal of Mathematics*, 85:351–390, 1994.

INSTITUTE OF MATHEMATICS THE HEBREW UNIVERSITY OF JERUSALEM JERUSALEM 91904, ISRAEL AND DEPARTMENT OF MATHEMATICS RUTGERS UNIVERSITY NEW BRUNSWICK, NJ 08854, USA

E-mail address: `shelah@math.huji.ac.il`